



# Epley versus Semont Maneuver in the Treatment of Benign Paroxysmal Positional Vertigo

Aravind Karuppan Muthu<sup>1</sup> Kiran Ravindranath Bongale<sup>1</sup> Vikram Kemmannu Bhat<sup>1</sup>

<sup>1</sup>Department of Otorhinolaryngology, Head and Neck Surgery, Karnataka Institute of Medical Sciences, Hubli, Karnataka, India

**Address for correspondence** Vikram Kemmannu Bhat, MBBS, MS, DNB, PhD, Department of Otorhinolaryngology, Head and Neck Surgery, Karnataka Institute of Medical Sciences, Vidyanagar, Hubli, 580021, Karnataka, India (e-mail: vikram.ent@gmail.com).

Ann Otol Neurotol ISO 2022;5:28–30.

## Abstract

### Keywords

- ▶ BPPV
- ▶ maneuver
- ▶ posterior semicircular canal

**Introduction** Benign paroxysmal positional vertigo (BPPV) is characterized by brief but violent attacks of paroxysmal vertigo provoked by certain positions of the head. This study aimed to compare the outcomes of Epley and Semont maneuvers for the relief of vertigo in posterior semicircular canal BPPV.

**Materials and Methods** This was a prospective comparative study conducted in a tertiary referral hospital over 1 year. Among 70 cases of BPPV, 35 were treated with Epley and 35 with Semont maneuver. Follow-up was done at 1 week, 1 month, and 3 months to assess symptomatic improvement using visual analogue scale.

**Results** Out of 70 patients of BPPV, 62 patients responded very well to the maneuvers and got relief from vertigo. Thirty-two patients (91.4%) in the Epley group and 30 (85.7%) patients in the Semont group had a significant improvement post-therapy and the balance was the same in both groups. At 1 month, there was no significant statistical difference in mean visual analogue score (VAS) score between both groups.

**Conclusion** Majority of cases showed symptomatic improvement with both maneuvers. However, patients treated with Epley maneuver showed earlier relief from vertigo than those treated with the Semont maneuver.

## Introduction

Benign paroxysmal positional vertigo (BPPV) is one of the common causes of peripheral vertigo with an estimated lifetime prevalence of 2.4%.<sup>1</sup> It may be associated with a reduced quality of life, falls, and depression. It is also the most rewarding vestibular condition to manage, which is characterized by brief attacks of vertigo associated with nystagmus precipitated by certain changes in the head position.

Posterior semicircular canal (PSC) BPPV has more incidence and prevalence than lateral and anterior semicircular canal BPPV.<sup>2</sup> The introduction of the procedures by Epley and Semont has revolutionized the management of BPPV to the point that most physicians, paramedical health care

providers, and even self-treated patients have used these maneuvers.<sup>3,4</sup> This study was conducted to assess the effectiveness of these maneuvers in the treatment of PSC BPPV.

## Materials and Methods

This non-randomized clinical study was undertaken on 70 patients in 1 year in a tertiary referral hospital after obtaining the necessary clearance from the institutional review board [KIMS/EC/66/2019–20]. Written informed consent was taken from all adult patients with PSC BPPV for the study. All of them underwent the Dix–Hallpike positional test and the diagnosis was confirmed with up beating torsional nystagmus. Cases with bilateral involvement,

DOI <https://doi.org/10.1055/s-0043-1761405>  
ISSN 2581-9607

© 2023. Indian Society of Otology.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Thieme Medical and Scientific Publishers Pvt. Ltd. A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

cervical spondylosis or neck injury, central vertigo, multicausal pathology were excluded from the study. Then they were allocated either Epley (35 patients) or the Semont maneuver (35 patients) as office procedure treatment and the outcome was graded based on symptomatic improvement. The treatment efficiency was evaluated by obtaining subjective relief of vertigo and was graded according to the visual analog score (VAS) from 0 to 10. After the maneuver, patient was instructed not to extend the head to the affected side and sleep with head elevated.

For the unresponsive patients, maneuvers were repeated after 1 week. Follow-up was done at the end of 1 week and 1 month. After 1 month, patient data were gathered by monthly telephone interviews to determine whether symptoms of position-induced vertigo had returned.

The data were analyzed with Stata version 14 using Chi square, analysis of variance (ANOVA), and independent *t* tests. A *p*-value less than 0.05 was considered statistically significant.

## Results

In this study, 70 (30 male and 40 female) patients of PSC BPPV were treated with either Epley or Semont maneuver. The study population was aged between 22 and 79 years (►Table 1). The mean age distribution was  $47.8 \pm 13.8$ . The mean age distribution of Epley and Semont groups was found to be 47.9 and 47.7 years, respectively. Right PSC involvement

was observed in 37 (52%) and left in 33 (47%). Two (2.8%) patients had associated headache, and eight (11%) had associated nausea and vomiting along with giddiness. The number of cases that recovered with either procedure from 1 week to 4 weeks is given in ►Table 2. The outcomes after either procedure at the end of 1 week and 1 month are given in ►Table 3. The VAS score analyzed using *t* test, and the change in the scores after treatment is given in ►Table 4.

At the end of first week, 88.57% in the Epley group and 82.85% in the Semont group had relief from vertigo. Out of 70 patients of PSC BPPV, 62 patients (88.57%) responded very well to the maneuvers and got relief from vertigo at the end of 1 month. Two patients had a recurrence of vertigo within the follow-up of 3 months.

## Discussion

Patients with vertiginous symptoms will ultimately be diagnosed with BPPV in 17 to 42% of cases, making BPPV the most frequent cause of vertigo.<sup>5,6</sup> The most widely used maneuver for the treatment of PSC BPPV is that of Epley.<sup>3</sup> This maneuver is a simple, effective way to treat PSC BPPV, the most common variant. Celis-Aguilar et al found this maneuver to be effective even for short-term treatment of BPPV.<sup>7</sup> A few authors opined that in resistant cases and/or cases with suspicion of cupulolithiasis, Semont liberatory maneuver may be used.<sup>4</sup> However, in our study we had more failures with Semont maneuver than with Epley. Cetin et al in their study found similar recovery and recurrence rates for both

**Table 1** Comparison of the age class interval in the groups

Age (y)	No. of cases Epley (%)	No. of cases Semont (%)
< 40	10 (28.6)	11 (31.4)
41–49	10 (28.6)	10 (28.6)
50–59	8 (22.9)	5 (14.3)
> 59	7 (20.0)	9 (25.7)
Total	35 (100)	35 (100)

**Table 2** Duration of vertigo in the patients

Duration	<i>n</i>	%
1 week	30	42.9
2 weeks	20	28.6
3–4 weeks	12	17.0
> 4 weeks	8	11.4
Total	70	100

**Table 3** Improvement of symptoms at the end of first week and first month

Symptoms	End of 1 week		End of 1 month	
	Epley (%)	Semont (%)	Epley (%)	Semont (%)
Improved	31 (88.57)	29 (82.85)	32 (91.4)	30 (85.7)
Not improved	4 (11.43)	6 (17.15)	3 (8.6)	5 (14.3)
Total	35 (100)	35 (100)	35 (100)	35 (100)
<i>p</i> -Value	0.52		0.452	

**Table 4** Change in VAS score with treatment

VAS score	Epley group Mean (SD)	Semont group Mean (SD)	<i>p</i> -Value
At baseline	10 (0)	10 (0)	1
After the treatment	5.1 (1.0)	5.7 (1.5)	0.027
After 1 week	2.7 (1.3)	3.6 (1.9)	0.028
After 1 month	1.5 (1.9)	1.9 (2.6)	0.466

Epley maneuver and Brandt–Daroff exercises.<sup>8</sup> However, Vijayaraj et al found that the Epley maneuver was superior and resulted in a positive impact on the quality of life as compared with Brandt–Daroff exercises.<sup>9</sup> Karanjai et al found healing rates of 87% in the Epley group, 75% in the Semont group, and 56% in the Brandt–Daroff group in a randomized controlled trial.<sup>10</sup> Acharya et al too found that Epley maneuver gave superior results than Semont in terms of remission of symptoms, but the same in preventing relapses.<sup>11</sup> Our study too agrees that Epley maneuver is superior to Semont, even though the difference was not statistically significant. Gupta et al found a significant difference between the procedures and the results revealed that 90, 73.33, and 50% patients improved with Epley, Semont, and Brandt–Daroff groups respectively.<sup>12</sup> Hence they concluded that Epley procedure was the best choice and Semont or Brandt–Daroff should be the least preferred in patients with PSC BPPV.<sup>12</sup> Sen et al too found that the Epley maneuver was more effective than the Semont in a randomized double blind study.<sup>13</sup> Gaur et al in a prospective study concluded that Epley maneuver with medicines was more effective than medicines alone.<sup>14</sup> Akula et al in a study compared Epley maneuver with reassurance among 72 patients and found a success rate of 93.75% and 6.25%, respectively, in the second follow up.<sup>15</sup> All these studies prove the sheer popularity of Epley maneuver due to superior results. Cohen et al in their study concluded that repositioning treatments were likely to be effective as long as the head was moved rapidly enough and through the correct planes in space.<sup>16</sup> Perez–Vazquez et al, in their review, found that more than 90% of patients could be successfully treated with maneuvers that move the particle back to the utricle.<sup>17</sup> Kader et al in their study found success rates of 90%, 85%, and 80% in Epley maneuver, rolling over maneuver and Brandt–Daroff exercises respectively.<sup>18</sup> Toupet et al compared Epley and Semont–Toupet maneuvers in their study and concluded that both had similar efficacy. Post maneuver restrictions did not modify the intensity of vertigo and dizziness in the observation period.<sup>19</sup> In our study too, post maneuver restrictions were similar in both the groups and did not make any difference.

## Conclusion

Epley maneuver was faster and significantly more effective than Semont in relieving the symptoms in PSC BPPV immediately after the maneuver and at the end of 1 week. Patients treated with the Epley maneuver showed earlier relief than patients treated with the Semont maneuver. However, this difference was not statistically significant at the end of 1 week and 1 month.

### Conflict of Interest

None declared.

## References

- 1 von Brevern M, Radtke A, Lezius F, et al. Epidemiology of benign paroxysmal positional vertigo: a population based study. *J Neurol Neurosurg Psychiatry* 2007;78(7):710–715
- 2 Whitney SL, Alghwiri A, Alghadir A. Physical therapy for persons with vestibular disorders. *Curr Opin Neurol* 2015;28(1):61–68
- 3 Epley JM. The canalith repositioning procedure: for treatment of benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg* 1992;107(3):399–404 10.1177/019459989210700310
- 4 Semont A, Freeys G, Vitte E. Curing the benign paroxysmal vertigo with a liberatory maneuver. *Adv Otolaryngol* 1988; 42 :290–293
- 5 Hanley K, O'Dowd T, Considine N. A systematic review of vertigo in primary care. *Br J Gen Pract* 2001;51(469):666–671
- 6 Katsarkas A. Benign paroxysmal positional vertigo (BPPV): idiopathic versus post-traumatic. *Acta Otolaryngol* 1999;119(7):745–749
- 7 Celis-Aguilar EM, Medina-Cabrera CA, Torrontegui-Zazueta LA, et al. Short-term effect of Epley maneuver as treatment for subjective benign paroxysmal positional vertigo. *Indian J Otolaryngol Head Neck Surg* 2022;74:545–549
- 8 Cetin YS, Ozmen OA, Demir UL, Kasapoglu F, Basut O, Coskun H. Comparison of the effectiveness of Brandt–Daroff vestibular training and Epley canalith repositioning maneuver in benign Paroxysmal positional vertigo long term result: a randomized prospective clinical trial. *Pak J Med Sci* 2018;34(3):558–563
- 9 Vijayaraj V. A comparison between the effect of Epley's manoeuvre and Brandt–Daroff exercise in improving the quality of life (QOL) in patients with benign paroxysmal positional vertigo (BPPV). *Int J Appl Dent Sci* 2018; 4 (2) :228–237
- 10 Karanjai S, Saha AK. Evaluation of vestibular exercises in the management of BPPV. *Indian J Otolaryngol Head Neck Surg* 2010;62(2):202–207
- 11 Acharya S, Biswal S, Dash S, Aparajitha A. Comparative study of Epley's and Semont's maneuver in treatment of benign paroxysmal positional vertigo in a tertiary care hospital. *Indian J Otol* 2020; 26 (4):211–214
- 12 Gupta AK, Sharma KG, Sharma P. Effect of Epley, Semont Maneuvers and Brandt–Daroff exercise on quality of life in patients with posterior semicircular canal benign paroxysmal positional vertigo (PSCBPPV). *Indian J Otolaryngol Head Neck Surg* 2019;71(1):99–103
- 13 Sen K, Sarkar A, Raghavan A. Comparative efficacy of Epley and Semont manoeuvre in benign paroxysmal positional vertigo: a prospective randomized double-blind study. *Astrocyte* 2016;3(2):96
- 14 Gaur S, Awasthi SK, Bhadouriya SK, Saxena R, Pathak VK, Bisht M. Efficacy of Epley's maneuver in treating BPPV patients: a prospective observational study. *Int J Otolaryngol* 2015;2015:487160
- 15 Akula S, Reddy LS, Kiran AS, Suresh AM. Clinical study of BPPV and the effectiveness of canalolith repositioning manoeuvre in subjects of BPPV. *Indian J Otolaryngol Head Neck Surg* 2022;74(1):96–102
- 16 Cohen HS, Sangi-Haghepeykar H. Canalith repositioning variations for benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg* 2010;143(3):405–412
- 17 Pérez-Vázquez P, Franco-Gutiérrez V. Treatment of benign paroxysmal positional vertigo. A clinical review. *J Otol* 2017;12(4):165–173
- 18 Kader HAA, Halim FNA, Rahman TTA. A comparative study on effectiveness of the rolling-over maneuver in rehabilitation of patients with posterior semicircular canal benign paroxysmal positional vertigo. *Egypt J Otolaryngol* 2014; 30 :88–93
- 19 Toupet M, Ferrary E, Bozorg Grayeli A. Effect of repositioning maneuver type and postmaneuver restrictions on vertigo and dizziness in benign positional paroxysmal vertigo. *ScientificWorldJournal* 2012;2012:162123